

# Glyphosate



Credit: Mnolf / Wikimedia Commons

*Roundup used in an apple orchard in Ciardes, Italy. Despite studies linking it to chronic diseases such as cancer, it is widely used not only in agriculture but in public areas such as parks.*



## Pesticide Action Network Asia and the Pacific

P.O. Box 1170

10850 Penang, Malaysia

Tel: (604) 657 0271 / 656 0381

Fax: (604) 658 3960

Email: panap@panap.net

Homepage: www.panap.net

Copyright ©Pesticide Action Network Asia and the Pacific. All rights reserved.

*Pesticide Action Network Asia and the Pacific (PAN AP) encourages the reproduction and use of this publication as long as PAN AP is properly acknowledged as the source and provided with a copy of the final work.*

**Meriel Watts, PhD**  
June 2012

## Trade names

Roundup, and many others.

## Uses

Broad spectrum (non-selective), systemic, post-emergence herbicide used to control annual and perennial plants including grasses, sedges, broadleaf weeds and woody plants.

It is used for crops, orchards, glasshouses, plantations, vineyards, pastures, lawns, parks, golf courses, forestry, roadsides, railway tracks, industrial areas, and home gardening.

Most used herbicide in the world.

## Regulatory status

### International

None.

### National

**Colombia:** In 2001, a court order stopped the government aerial spraying of Roundup on illegal coca plantations on the border of Columbia and Ecuador (Antoniou et al 2010).

**Israel:** A court order stopped aerial spraying of Roundup on Bedouin farmers' crops in the Negev region of Israel between 2002 and 2004, after a coalition of Arab human rights groups and Israeli scientists reported high death rates of livestock, and a high incidence of miscarriages and disease amongst the people (Antoniou et al 2010).

**Argentina:** In 2010, a regional court in Argentina banned the spraying of glyphosate near populated areas of Santa Fe province. In 2012, a network of 160 physicians, health workers and researchers demanded a ban on aerial spraying of glyphosate and other pesticides in Argentina, based on increases in cancer and a range of other illnesses since the introduction of glyphosate-tolerant GM soybeans. The illnesses affect development, reproduction, and skin; and immune, respiratory, neurological, and endocrine systems (Robinson 2010; Sirinathsinghji 2012).

## Manufacture

Monsanto, and many other companies. Global production capacity is 1.1 million tonnes, and global demand is about 0.5 million tonnes. Most comes from China – its capacity is 0.8 million tonnes and it exports 0.3 million tonnes.

## Residues in food

Residues analysis is expensive and not often carried out, but residues have been found in bread, flour, wheat, barley, bran, oats, breakfast cereals, cereal bars, polenta strawberries, lettuce, carrots, soy, wild berries, and drinking water.

## Health effects

### Poisonings

Glyphosate herbicides have been frequently used in self-poisonings with many deaths, especially in Asia. Numerous occupational and bystander exposures have been reported.

In Columbia and Ecuador, widespread human poisonings have been reported following aerial spraying of cocoa crops with glyphosate. Effects included skin problems, abdominal pain, gastrointestinal problems, acute respiratory infection, conjunctivitis, fever, allergies, kidney disease, lupus, cancer and birth defects. Many fish, cattle and poultry died (Oldham & Massey 2002; Trigona 2009; Valente 2009).

In 2003, 11 year-old Silvino Talavera died after direct exposure to pesticides used on soybean fields. His mother and siblings were hospitalised for 3 months. In 2004, a court convicted 2 men of culpable homicide caused by the irresponsible and criminal use of agrochemicals sprayed on soybean, specifically glyphosate. Three family members had glyphosate residues in their bodies (Williamson 2004).

### Residues in people

Residues were found in all samples of urine from people in Berlin, ranging from 0.5 to 2.0 mg/L, assumed to come from food especially crops desiccated

pre-harvest by glyphosate (Brändli & Reinacher 2012).

Residues have also been found in the urine of farmers using glyphosate at level of 0.23 mg/L (Acquavella et al 2004).

### **Acute toxicity**

Signs and symptoms of exposure include:

- irritation, swelling, tingling, itching or burning of the skin, photo-contact dermatitis, recurrent eczema, blisters, rashes;
- numbness in the face, swelling of the eye and lid, face, and joints;
- conjunctivitis, painful eyes, corneal injury, burning eyes, blurred vision, weeping eyes;
- oral and nasal discomfort, unpleasant taste, tingling and irritation of throat, sore throat;
- difficulty breathing, cough, coughing of blood, inflammation of lungs;
- nausea, vomiting, headache, fever, diarrhoea, weakness;
- rapid heartbeat, palpitations, raised blood pressure, dizziness, chest pains (IPCS 1994; Cox 1998; Gallardo 2001; Bradberry et al 2004).

Dr Ricky Gorringer of New Zealand estimated, based on cases presenting to his clinic, that 1 in 20 New Zealanders are sensitive to Roundup. The most commonly occurring symptoms are unnatural fatigue, band-like headache, “spaced-out feeling with loss of confidence”, skin rash, and an otherwise unexplainable sudden increase in blood pressure. These problems are thought to largely result from micro-droplet inhalation (Watts 1994).

“Widespread disturbances of many body systems . . . after exposures at normal use levels. These include balance disorder, vertigo, reduced cognitive capacity, seizures, impaired vision, smell, hearing and taste, headaches, drops in blood pressure, body-wide twitches and tics, muscle paralysis, peripheral neuropathy, loss of gross and fine motor skills, excessive sweating and severe fatigue” (Ho & Ching 2003).

### **Chronic toxicity**

Formulated products contain so-called ‘inert’ ingredients in addition to the active ingredient. Sometimes they can be more toxic than the active, and this is particularly so for Roundup, hence studies are often carried out separately on Roundup and glyphosate.

**General:** decreased body weight gain, increased incidence of cataract and lens abnormalities, increased liver weight, and degeneration of the liver and kidney at high doses in laboratory trials. Concentrations far below those normally sprayed kill human skin cells, cause aging of the skin, and can lead to skin cancer. (US EPA 1993; Heu et al 2012 a,b)

**Genotoxicity:** glyphosate and/or Roundup are genotoxic in human lymphocytes, liver cells, and mouth cells; and in mouse bone marrow, liver and kidney cells, bovine lymphocytes, fruit flies, fish, alligators, tadpoles, sea urchins, onion root-tip cells, and bacteria. Epidemiological studies show DNA damage in people exposed to aerial spraying of glyphosate in Ecuador and Colombia. One of the main health implications of genotoxicity is cancer. (Vigfusson & Vyse 1980; Rank et al 1993; Kale et al 1995; Bolognesi et al 1997, 2009; Clements et al 1997, Lioi et al 1998a,b; Peluso et al 1998; Kaya et al 2000; Grisolia 2002; Mueckay & Maldonado 2003; Monroy et al 2005; Siviková & Dianovský 2006; Bellé et al 2007; Çavaş & Könen 2007; Cavalcante et al 2008; Gasnier et al 2009; Guilherme et al 2009; Mañas et al 2009a,b; Poletta et al 2009; Prasad et al 2009; Koller et al 2012)

**Other cancer-causing mechanisms:** glyphosate or Roundup cause cell division dysfunction, cell-cycle dysregulation at concentrations hundreds of times lower than those sprayed. They inhibit RNA transcription at concentrations 25 times lower than those sprayed; and cause oxidative stress in human lymphocytes and skin cells, bovine lymphocytes, bullfrog tadpoles, pregnant rats and their foetuses, rat liver cells, mouse kidney cells and liver DNA, fish brain, liver and kidney cells, and in rice leaves. Additionally glyphosate enhanced the genotoxic effect of hydrogen peroxide through an oxidative stress mechanism. Roundup caused mild oxidative stress in the brain, liver and kidneys of goldfish. (Lioi et al 1998a; Lueken et al 2004; Pieniżek et al 2004; Gehin et al 2005, 2006; Lioi et al 1998b; Costa et al 2008; Beuret et al 2005; El-Shenawy 2009; Bolognesi et al 1997; Marc et al 2002, 2003, 2004, 2005; Ahsan et al 2008; Lushchak et al 2009)

**Cancer:** thyroid C-cell carcinomas, liver carcinoma, leukaemia, lymphoma; adenomas of pancreas, liver, thyroid in rodents at high doses. Studies on mice show it has

tumour-promoting potential for skin cancer. Both Roundup and glyphosate cause proliferation of breast cancer cells. (US EPA 1993; Hardell & Eriksson 1999; Lin & Garry 2000; Hokanson et al 2007; George et al 2009)

Epidemiological studies indicate non-Hodgkin’s lymphoma and multiple myeloma. Doctors and the state government of Chaco report increased incidence of cancer in Argentina following the introduction of GM soy crops that are heavily sprayed with Roundup. (McDuffie et al 2001; Hardell et al 2002; De Roos et al 2003; Robinson 2010; Chaco 2010)

**Endocrine disruption:** Roundup but not glyphosate inhibits the conversion of androgens to oestrogen, disrupts the production of progesterone, and reduces testosterone levels. Both Roundup and glyphosate dysregulate oestrogen-dependent breast cancer cells, and interfere with aromatase, the enzyme that converts androgens to oestrogen. Glyphosate is anti-androgenic at concentrations 40 times lower than residues permitted in GM soybeans. (Walsh et al 2000; Gasnier et al 2010; Hokanson et al 2007; Clair et al 2012a).

**Reproductive and developmental toxicity:** Monsanto has known since 1993 that low doses of glyphosate cause malformations in animals. These include reduced ossification and higher incidence of skeletal and organ anomalies; and reduced number of viable foetuses (Antoniu et al 2011).

Rats treated with sublethal doses of Roundup Max during pregnancy had decreased implantation rates, increased resorption of foetuses, and the surviving offspring had decreased skeletal calcification (Gerişlioğlu et al 2010).

Roundup Transorb significantly altered the progression of puberty, reduced testosterone production and altered seminiferous tubules (Romano et al 2010).

Glyphosate, Roundup, the adjuvant POEA, and the metabolite AMPA all cause cell death in human umbilical, embryonic and placental cells, at dilutions far below those used in agriculture. These effects together with the endocrine disrupting effects can result in pregnancy problems leading to abnormal foetal development, low birth weights, or miscarriages. Glyphosate kills testicular Sertoli

cells. (Benachour et al 2007; Benachour & Seralini 2009; Gammon 2009; Clair et al 2012a)

Birth defects are reported in Argentina and Paraguay amongst those exposed to aerial spraying of glyphosate. These are neural tube defects (absence of the majority of the brain, skull and scalp), small head, and craniofacial malformations, strikingly similar to those caused by glyphosate in laboratory studies on amphibia.

There has been a 4-fold increase in birth defects in the Argentinean state of Chaco with the introduction of GM Roundup Ready crops (Benítez-Leite et al 2009; Paganelli et al 2010; Antoniou et al 2010; Chaco 2010). Epidemiological studies have reported miscarriages, pre-term deliveries, and reduced fecundity (Savitz et al 1997; Curtis et al 1999; Arbuckle et al 2001).

**Neurotoxicity:** Glyphosate inhibits the growth of axons and dendrites in nerve cells. It also depletes the neurotransmitters serotonin and dopamine, and damages the substantia nigra region of the brain, implicating them in Parkinson's disease (Axelrad et al 2003; Anadón et al 2008; Astiz et al 2009).

**Immunotoxicity:** exposure to glyphosate has been associated with allergic responses; one case developed autoimmune blistering of skin and mucous membranes after exposure to the fumes of burning glyphosate. Immune effects have been found in fish. Immune function effects have been reported from Argentina in areas of heavy aerial spraying of glyphosate over GM crops. (el-Gendy et al 1998; Slager et al 2009; Fisher et al 2008; Sirinathsinghji 2012).

## Environmental and agroecological effects

### Toxicity

**Aquatic:** Glyphosate is increasingly found in the aquatic environment at levels that have caused significant effects on species that underpin the entire aquatic food chain. It can alter the composition of natural aquatic communities, tipping the ecological balance, changing clear water to turbid water, giving rise to harmful algal blooms, and reducing species richness. It can have profound impacts on microorganisms, plankton, algae and amphibia at low concentrations: one study showed a 70% reduction

in tadpole species and a 40% increase in algae. Insects, molluscs, crustaceans, sea urchins, reptiles, tadpoles, and fish can all be affected, vulnerability within each group varying dramatically between species. Effects include reproductive abnormalities, developmental abnormalities and malformations, DNA damage, immune effects, oxidative stress, modified enzyme activity, decreased capacity to cope with stress and maintain homeostasis, altered behaviour, and impaired olfaction that can threaten their survival. Amphibians are particularly vulnerable.

Roundup is generally more toxic than glyphosate, especially to fish. (Relyea 2005; Pérez et al 2007; Vera et al 2009; Saxton et al 2011).

**Terrestrial:** Glyphosate at concentrations lower than those used in agriculture inhibited the growth of 3 microorganisms important in food processing. It interferes with key molecular mechanisms which regulate early development in chickens leading to congenital malformations (Paganelli et al 2010; Clair et al 2012b).

### Agroecological disruption

**Beneficials:** Glyphosate has adverse effects on some earthworms; and on a number of beneficial insects useful in biological control, particularly predatory mites, carabid beetles, ladybugs, spiders, and green lacewings. It can also adversely affect other insects that play an important part in ecological balance such as springtails, wood louse, and field spiders (Schneider et al 2009; Benamú et al 2010; Griesinger et al 2011).

**Soil organisms:** Glyphosate has adverse effects on some earthworms. Glyphosate is toxic to some but not all soil microorganisms, altering microbial community dynamics in ways that are harmful to plants and to ecological balance. It increases microorganisms capable of metabolising the chemical. It can reduce some beneficial organisms such as saprophytic fungi that decompose dead plant material and are important for soil fertility. Glyphosate is toxic to nitrogen-fixing bacteria, and also increases denitrification contributing to nitrous oxide emissions and nitrogen loss from soil. (Grossbard 1985; Carlisle & Trevors 1988; Tenuta & Beauchamp 1996; Zablutowicz & Reddy 2007; Kremer & Means 2009; Lupwayi et al 2009).

**Crop effects:** Numerous studies show glyphosate stimulates growth of some fungal pathogens, such as *Fusarium*, that cause diseases in many crops. The upsurge in use of glyphosate in no-till agriculture has brought about a resurgence of some diseases. Glyphosate binds micronutrients in the soil and causes micronutrient deficiencies in plants that increase their susceptibility to disease, decrease their vigour, and produce micronutrient-deficient food crops. It can reduce the plant's production of lignin and phenolic compounds, which are also important for disease resistance. It can reduce nitrogen-fixation in legumes such as soybean. It causes nutritional changes in soy including reduced iron, linoleic acid and linolenic acid, and increased monounsaturated fatty acids (Ozturk et al 2008; Johal & Huber 2009; Zobiole et al 2010).

**Resistance:** 23 species of plants in 20 countries are resistant to glyphosate and Monsanto is encouraging farmers to use dicamba or 2,4-D alongside the Roundup (Monsanto 2012; Schiffman 2012; WeedScience.com 2012).

### Environmental fate and contamination

**Air and rain:** Glyphosate and the metabolite AMPA were found in up to 100% of samples of rain and air in agricultural areas in the US; about 0.7% of glyphosate applied in agricultural areas is removed from the air in rain. It was one of the most frequently detected pesticides in rainwater in Belgium in 2001; also measured in rain in Canada, at all sites and throughout the "growing season", at a maximum concentration of 1.51 ug/m<sup>2</sup>/day (Quaghebeur et al 2004; Humphries et al 2005; Chang et al 2011).

**Soil:** relatively persistent (up to 180 days in some soils). The metabolite AMPA accumulates: 20 years after application, 8.5% of the amount applied is likely to remain in soils (Mamy et al 2010).

**Aquatic:** Runoff from hard surfaces in urban areas is increasingly seen as a source of surface water pollution with glyphosate and AMPA. Residues of glyphosate and AMPA were found in surface water in Argentina, Canada, China, France, Netherlands, Norway, Switzerland, Germany, UK, USA; in wastewater in France and Canada, landfill leachate in the UK, and groundwater in Spain. (Slack et al

2005; Ghanem et al 2007; Peruzzo et al 2008; Hanke et al 2010; Mamy et al 2010; Székács & Davas 2011)

Glyphosate has been found in groundwater and wells in a number of countries, including Canada, Denmark, the Netherlands, and USA (IPCS 1994, US EPA 1992; Cox 1998).

Both glyphosate and AMPA have been found in marine sediment in the Waitemata Harbour and Hauraki Gulf in New Zealand, believed to have come largely from the spraying of urban roadside weeds. The maximum glyphosate concentration detected was 1 ppm, while AMPA had a maximum level of 0.37 ppm (Stewart et al 2009).

**Bioaccumulation:** Low-level bioaccumulation has been found in snails, fish, and freshwater worm (Wang et al 1994; Contardo-Jara et al 2009; Druart et al 2011).

## Sources include

Acquavella JF, Alexander BH, Mandel JS, Gustin C, Baker B, Chapman P, Bleeke M. 2004. Glyphosate biomonitoring for farmers and their families: results from the farm family exposure study. *Environ Health Perspect* 112(3):321-6.

Ahsan N, Lee DG, Lee KW, Alam I, Lee SH, Bahk JD, Lee BH. 2008. Glyphosate-induced oxidative stress in rice leaves revealed by proteomic approach. *Plant Physiol Biochem* 46(12):1062-70.

Anadón A, del Pino J, Martínez MA, Caballero V, Ares I, Nieto I, Martínez-Larrañaga MR. 2008. Neurotoxicological effects of the herbicide glyphosate. *Toxicol Lett* 180S:S164.

Antoniou M, Brack P, Carrasco A, Fagan J, Habib M, Kageyama P, Leifert C, Nodari RO, Pengue W. 2010. GM Soy Sustainable? Responsible? GLS Bank, Germany. [http://www.global2000.at/module/media/data/global2000\\_at\\_de/content/globalnews/carrasco\\_etal\\_gmsoy\\_respons\\_full\\_study.pdf\\_me/carrasco\\_etal\\_gmsoy\\_respons\\_full\\_study.pdf](http://www.global2000.at/module/media/data/global2000_at_de/content/globalnews/carrasco_etal_gmsoy_respons_full_study.pdf_me/carrasco_etal_gmsoy_respons_full_study.pdf).

Antoniou M, Habib M, Howard CV, Jennings RC, Leifert C, Nodari RO, Robinson C, Fagan J. 2011. Roundup and Birth Defects: Is the Public Being Kept in the Dark? Earth Open Source. <http://www.scribd.com/doc/57277946/RoundupandBirthDefects5v5>.

Arbuckle TE, Lin Z, Mery LS. 2001. An exploratory analysis of the effect of pesticide exposure on the risk of spontaneous abortion in an Ontario farm population. *Environ Health Perspect* 109:851-57.

Astiz M, de Alaniz MJT, Marra CA. 2009. Effect of pesticides on cell survival in liver and brain rat tissues. *Ecotoxicol Environ Saf* 72(7):2025-32.

Axelrad JC, Howard CV, McLean WG. 2003. The effects of acute pesticide exposure on neuroblastoma cells chronically exposed to diazinon. *Toxicology* 185:67-78.

Bellé R, Le Bouffant R, Morales J, Cosson B, Cormier P, Mulner-Lorillon O. 2007. [Sea urchin embryo, DNA-damaged cell cycle checkpoint and the mechanisms initiating cancer development]. *J Soc Biol* 201(3):317-27.

Benachour N, Sipahutar H, Moslemi S, Gasnier C, Travert C, Séralini G-E. 2007. Time- and dose-dependent effects of Roundup on human embryonic and placental cells. *Arch Environ Contam Toxicol* 53:126-33.

Benachour N, Séralini G-É. 2009. Glyphosate formulations induce apoptosis and necrosis in human umbilical, embryonic, and placental cells. *Chem Res Toxicol* 22:97-105.

Benamú MA, Schneider MI, Sánchez NE. 2010. Effects of the herbicide glyphosate on biological attributes of *Alpaida veniliae* (Araneae, Araneidae), in laboratory. *Chemosphere* 78:871-6.

Benítez-Leite S, Macchi MA, Acosta M. 2009. Malformaciones congénitas asociadas a agrotóxicos. *Arch Pediatr Urug* 80:237-47. Cited in Paganelli et al 2010.

Beuret CJ, Zirulnik F, Giménez MS. 2005. Effect of the herbicide glyphosate on liver lipoperoxidation in pregnant rats and their fetuses. *Repro Toxicol* 19(4):501-4.

Brändli D, Reinacher S. 2012. Herbicides found in human urine. *Ithaka Journal* Jan 270-272.

Bolognesi C, Bonatti S, Degan P, Gallerani E, Peluso M, Rabboni R, Roggieri P, Abbondandolo A. 1997. Genotoxic activity of glyphosate and its technical formulation Roundup. *J Agric Food Chem* 45(5):1957-62.

Bolognesi C, Carrasquilla G, Volpi S, Solomon KR, Marshall EJP. 2009. Biomonitoring of genotoxic risk in agricultural workers from five Colombian regions: association to occupational exposure to glyphosate. *J Toxicol Environ Health A* 72(15&16):986-97.

Bradberry SM, Proudfoot AT, Vale JA. 2004. Glyphosate poisoning. *Toxicol Rev* 23(3):159-67.

Carlisle SM, Trevors JT. 1988. Glyphosate in the environment. *Water Air Soil Pollut* 39 (3-4):409-20.

Cavalcante DGSM, Martínez CBR, Sofia SH. 2008. Genotoxic effects of Roundup on the fish *Prochilodus lineatus*. *Mutat Res* 655(1-2):41-6.

Çavaş T, Könen S. 2007. Detection of cytogenetic and DNA damage in peripheral erythrocytes of goldfish (*Carassius auratus*) exposed to a glyphosate formulation using the micronucleus test and the comet assay. *Mutagenesis* 22(4):263-8.

Chaco. 2010. Primer Informe de Comisión Provincial de Investigación de Contaminantes del Agua. [http://www.gmwatch.eu/files/Chaco\\_Government\\_Report\\_Spanish.pdf](http://www.gmwatch.eu/files/Chaco_Government_Report_Spanish.pdf).

Chang F-C, Simcik MF, Capel PD. 2011. Occurrence and fate of the herbicide glyphosate and its degradate aminomethylphosphonic acid in the atmosphere. *Environ Toxicol Chem* 30(3):548-55.

Clair E, Mesnage R, Travert C, Séralini G-E. 2012a. A glyphosate-based herbicide induces necrosis and apoptosis in mature rat testicular cells in vitro, and testosterone decrease at lower levels. *Toxicol In Vitro* 26:269-79.

Clair E, Linn L, Travert C, Amiel C, Séralini G-É, Panoff J-M. 2012b. Effects of Roundup and glyphosate on three food microorganisms: *Geotrichum candidum*, *Lactococcus lactis* subsp. *cremoris* and *Lactobacillus delbrueckii* subsp. *bulgaricus*. *Curr Microbiol* 64(5):486-91.

Clements C, Ralph S, Petras M. 1997. Genotoxicity of select herbicides in *Rana catesbeiana* tadpoles using the alkaline single-cell gel DNA electrophoresis (comet) assay. *Environ Mol Mutagen* 29(3):277-88.

Contardo-Jara V, Klingelmann E, Wiegand C. 2009. Bioaccumulation of glyphosate and its formulation Roundup Ultra in *Lumbriculus variegatus* and its effects on biotransformation and antioxidant enzymes. *Environ Pollut* 157:57-63.

Costa MJ, Monteiro DA, Oliveira-Neto AL, Rantin FT, Kalinin AL. 2008. Oxidative stress biomarkers and heart function in bullfrog tadpoles exposed to Roundup Original. *Ecotoxicol* 17(3):153-63.

Cox C. 1998. Glyphosate (Roundup). *J Pestic Reform* 18(3):3-16.

Curtis KM, Savitz DA, Weinberg CR, Arbuckle TE. 1999. The effect of pesticide exposure on time to pregnancy. *Epidemiology* 10(2):112-7.

De Roos AJ, Zahm SH, Cantor KP, Weisenburger DD, Holmes FF, Burmeister LF, Blair A. 2003. Integrative assessment of multiple pesticides as risk factors for non-Hodgkin's lymphoma among men. *Occup Environ Med* 60:E11.

Druart C, Millet M, Scheifer R, Delhomme O, Raappel C, de Vaulfleury A. 2011. Snails as indicators of pesticide drift, deposit, transfer and effects in the vineyard. *Sci Total Environ* 409(20):4280-8.

el-Gendy KS, Aly NM, el-Sebae AH. 1998. Effects of edifenphos and glyphosate on the immune response and protein biosynthesis of boliti fish (*Tilapia nilotica*). *J Environ Sci Health B* 33(2):135-49.

El-Shenawy NS. 2009. Oxidative stress responses of rats exposed to Roundup and its active ingredient glyphosate. *Environ Toxicol Pharmacol* 28(3):379-85.

Fisher KR, Higginbotham R, Frey J, Granese J, Pillow J, Skinner RB. 2008. Pesticide-associated pemphigus vulgaris. *Cutis* 82(1):51-4.

Gallardo L. 2001. Aerial herbicide impact on farmers in Ecuador. *Pestic News* 54:8.

Gammon C. 2009. Weed killer kills human cells. Study intensifies debate over 'inert' ingredients. *Environ Health News* June 22. <http://www.environmentalhealthnews.org/ehs/news/roundup-weed-killer-is-toxic-to-human-cells-study-intensifies-debate-over-inert-ingredients>.

Gasnier C, Dumont C, Benachour N, Clair E, Chagnon M-C, Séralini G-E. 2009. Glyphosate-based herbicides are toxic and endocrine disruptors in human cell lines. *Toxicology* 262(3):184-91.

Gasnier C, Benachour N, Clair E, Travert C, Langlois F, Laurant C, Decroix-Laporte C, Séralini G-E. 2010. Dig1 protects against cell death provoked by glyphosate-based herbicides in human liver cell lines. *J Occup Med Toxicol* 27:5-29.

Gehin A, Guillaume YC, Millet J, Guyon C, Nicod L. 2005. Vitamins C and E reverse effect of herbicide-induced toxicity on human epidermal cells HaCaT: a biochemometric approach. *Int J Pharm* 288:219-26.

Gehin A, Guyon C, Nicod L. 2006. Glyphosate-induced antioxidant imbalance in HaCaT: the protective effect of vitamins C and E. *Environ Toxicol Pharmacol* 22:27-34.

George J, Prasad S, Mahmood Z, Shukla Y. 2009. Studies on glyphosate-induced carcinogenicity in mouse skin: a proteomic approach. *J Proteomics* 73:951-64.

Gerişlioğlu A, Güngörmüş C, Korkmaz A, Kolankaya D. 2010. Embryotoxic and teratogenic effects of Roundup max on rat development. *Toxicol Lett* 196S(S37-351):P204-005.

Ghanem A, Bados P, Estaun AR, de Alencastro LF, Taibi S, Einhorn J, Mouglin C. 2007. Concentrations and specific loads of glyphosate, diuron, atrazine, nonylphenol and metabolites thereof in French urban sewage sludge. *Chemosphere* 69:1368-73.

Griesinger LM, Evans SC, Rypstra AL. 2011. Effects of a glyphosate-based herbicide on mate location in wolf spider that inhabits agroecosystems. *Chemosphere* 84:1461-6.

Grisolia CK. 2002. A comparison between mouse and fish micronucleus test using cyclophosphamide, mitomycin C and various pesticides. *Mutat Res* 518:145-50.

Grossbard E. 1985. Effects of glyphosate on the microflora: with reference to the decomposition of treated vegetation and interaction with some plant pathogens. In: Grossbard E, Atkinson D (eds). *The Herbicide Glyphosate*. Butterworths, London.

Guilherme S, Gaivão I, Santos MA, Pacheco M. 2009. Tissue specific DNA damage in the European eel (*Anguilla anguilla*) following a short-term exposure to a glyphosate-based herbicide. *Toxicol Lett* 189S:S212:Z15.

Hanke I, Witterer I, Bischofberger S, Stamm C, Singer H. 2010. Relevance of urban glyphosate use for surface water quality. *Chemosphere* 81:422-9.

Hardell L, Eriksson M. 1999. A case-control study of non-Hodgkin lymphoma and exposure to pesticides. *Cancer* 85(6):1353-60.

Hardell L, Eriksson M, Nordstrom M. 2002. Exposure to pesticides as risk factor for non-Hodgkin's lymphoma and hairy cell leukemia: pooled analysis of two Swedish case-control studies. *Leuk Lymphoma* 43:1043-9.

Heu C, Elie-Caille C, Mougey V, Launay S, Nicod L. 2012a. A step further toward glyphosate-induced epidermal cell death: Involvement of mitochondrial and oxidative mechanisms. *Environ Toxicol Pharmacol* 34:144-53.

Heu C, Berquand A, Elie-Caille C, Nicod L. 2012b. Glyphosate-induced stiffening of HaCaT keratinocytes, a Peak Force Tapping study in living cells. *J Struct Biol* 178:1-7.

Ho M-W, Ching LL. 2003. The Case For A GM-Free Sustainable World. Independent Science Panel, London. <http://www.i-sis.org.uk>

Hokanson R, Fudge R, Chowdhary R, Busbee D. 2007. Alteration of estrogen-regulated gene expression in human cells induced by the agricultural and horticultural herbicide glyphosate. *Hum Exp Toxicol* 26:747-52.

Humphries D, Byrtus G, Anderson A-M. 2005. Glyphosate Residues in Alberta's Atmospheric Deposition, Soils, and Surface Waters. Pub No. T/806, Alberta Environment, Edmonton. <http://environment.gov.ab.ca/info/library/6444.pdf>.

- IPCS. 1994. Environmental Health Criteria 159: Glyphosate. International Programme on Chemical Safety, World Health Organisation, Geneva. <http://www.inchem.org/documents/ehc/ehc/ehc159.htm>.
- Johal GS, Huber DM. 2009. Glyphosate effects on diseases of plants. *Eur J Agron* 31(3):144-52.
- Kale PG, Petty BT Jr, Walker S, Ford JB, Dehkordi N, Tarasia S, Tasić BO, Kale R, Sohni YR. 1995. Mutagenicity testing of nine herbicides and pesticides currently used in agriculture. *Environ Mol Mutagen* 25:148-53.
- Kaya B, Creus A, Yanikoğlu A, Cabré O, Marcos R. 2000. Use of the *Drosophila* wing spot test in the genotoxicity testing of different herbicides. *Environ Mol Mutagen* 36(1):40-6.
- Koller VJ, Fühacker M, Nersisyan A, Mišák M, Eisenbauer M, Knasmueller S. 2012. Cytotoxic and DNA-damaging properties of glyphosate and Roundup in human-derived buccal epithelial cells. *Arch Toxicol* 86(5):805-13.
- Kremer RJ, Means NE. 2009. Glyphosate and glyphosate-resistant crop interactions with rhizosphere microorganisms. *Eur J Agron* 31(3):153-61.
- Lin N, Garry VF. 2000. In vitro studies of cellular and molecular development toxicity of adjuvants, herbicides, and fungicides commonly used in Red River Valley, Minnesota. *J Toxicol Environ Health A* 60:423-39.
- Lioi MB, Scarfi MR, Santoro A, Barbieri R, Zeni O, Salvemini F, Di Berardino D, Ursini MV. 1998a. Cytogenetic damage and induction of pro-oxidant state in human lymphocytes exposed in vitro to glyphosate, vinclozolin, atrazine, and DPX-E9636. *Environ Mol Mutagen* 32:39-46.
- Lioi MB, Scarfi MR, Santoro A, Barbieri R, Zeni O, Di Berardino D, Ursini MV. 1998b. Genotoxicity and oxidative stress induced by pesticide exposure in bovine lymphocyte cultures in vitro. *Mutat Res* 403(1-2):13-20.
- Lueken A, Juhl-Strauss U, Krieger G, Witte I. 2004. Synergistic DNA damage by oxidative stress (induced by H<sub>2</sub>O<sub>2</sub>) and nongenotoxic environmental chemicals in human fibroblasts. *Toxicol Lett* 147(1):35-43.
- Lupwayi NZ, Harker KN, Clayton GW, O'Donovan JT, Blackshaw RE. 2009. Soil microbial response to herbicides applied to glyphosate-resistant canola. *Agric Ecosys Environ* 129(1-3):171-6.
- Lushchak OV, Kubrak OI, Storey JM, Storey KB, Lushchak VI. 2009. Low toxic herbicide Roundup induces mild oxidative stress in goldfish tissues. *Chemosphere* 76(7):932-7.
- Mamy L, Gabrielle B, Barriuso E. 2010. Comparative environmental impacts of glyphosate and conventional herbicides when used with glyphosate-tolerant and non-tolerant crops. *Environ Pollut* 158:3172-8.
- Mañas F, Peralta L, Raviolo J, García Ovando H, Weyers A, Ugnia L, Gonzalez Cid M, Larripa I, Gorla N. 2009a. Genotoxicity of glyphosate assessed by the comet assay and cytogenetic tests. *Environ Toxicol Pharmacol* 28:37-41.
- Mañas F, Peralta L, Raviolo J, García Ovando H, Weyers A, Ugnia L, Gonzalez Cid M, Larripa I, Gorla N. 2009b. Genotoxicity of AMPA, the environmental metabolite of glyphosate, assessed by the Comet assay and cytogenetic tests. *Ecotoxicol Environ Saf* 72(3):834-7.
- Marc J, Mulner-Lorillon O, Boulben S, Hureau D, Durand G, Bellé R. 2002. Pesticide Roundup provokes cell division dysfunction at the level of CDK1/cyclin B activation. *Chem Res Toxicol* 15(3):326-31.
- Marc J, Mulner-Lorillon O, Durand G, Bellé R. 2003. Embryonic cell cycle for risk assessment of pesticides at the molecular level. *Environ Chem Lett* 1(1):8-12.
- Marc J, Mulner-Lorillon O, Bellé R. 2004. Glyphosate-based pesticides affect cell cycle regulation. *Biol Cell* 96(3):245-9.
- Marc J, Le Breton M, Cormier P, Morales J, Bellé R, Mulner-Lorillon O. 2005. A glyphosate-based pesticide impinges on transcription. *Toxicol Appl Pharmacol* 203:1-8.
- McDuffie HH, Pahwa P, McLaughlin JR, Spinelli JJ, Fincham S, Dosman JA, Robson D, Skinnider LF, Choi NW. 2001. Non-Hodgkin's lymphoma and specific pesticide exposures in men: cross-Canada study of pesticides and health. *Cancer Epidemiol Biomarkers Prev* 10:1155-63.
- Monroy CM, Cortés AC, Sicard DM, de Restrepo HG. 2005. [Cytotoxicity and genotoxicity of human cells exposed in vitro to glyphosate]. *Biomedica* 25(3):335-45.
- Monsanto. 2012. New Roundup® Ready Xtend Crop System to Extend Weed Control and Maximize Yield. Monsanto Company, St Louis. <http://cornandsoybeandigest.com/new-roundup-ready-xtend-crop-system-extend-weed-control-and-maximize-yield>.
- Mueckay C, Maldonado A. 2003. Daños genéticos en la frontera de Ecuador por las fumigaciones del Plan Colombia. Acción Ecológica, Acción Creativa, Clínica de Derechos Humanos de la PUCE, CAS, CEDHU, CONAIE, FORCCOFES, INREDH, Plan Pais, RAPAL Ecuador, SERPAJ Ecuador. <http://www.visionesalternativas.com/militarizacion/articulos/peolom/AE0311.pdf>.
- Oldham J, Massey R. 2002. Health and Environmental Effects of Herbicide Spray Campaigns in Colombia. Institute for Science and Interdisciplinary Studies, Amherst MA. [http://laslanas.org/Colombia/SprayingReview\\_Oldham-Massey.pdf](http://laslanas.org/Colombia/SprayingReview_Oldham-Massey.pdf).
- Ozturk L, Yacizi A, Eker S, Gokmen O, Römheld V, Cakmak I. 2008. Glyphosate inhibition of ferric reductase activity in iron deficient sunflower roots. *New Phytol* 177(4):899-906.
- Paganelli A, Gnazzo V, Acosta H, López SL, Carrasco AE. 2010. Glyphosate-based herbicides produce teratogenic effects on vertebrates by impairing retinoic acid signalling. *Chem Res Toxicol* [Epub May].
- Peluso M, Munni A, Bolognesi C, Parodi S. 1998. 32P-postlabeling detection of DNA adducts in mice treated with the herbicide Roundup. *Environ Mol Mutagen* 31:55-9.
- Pérez GL, Torremorell A, Mugni H, Rodríguez P, Solange Vera M, do Nascimento M, Allende L, Bustingorry J, Escaray R, Ferraro M, Izaguirre I, Pizarro H, Bonetto C, Morris DP, Zagarese H. 2007. Effects of the herbicide Roundup on freshwater microbial communities: a mesocosm study. *Ecol Appl* 17(8):2310-22.
- Peruzzo PJ, Porta AA, Ronco AE. 2008. Levels of glyphosate in surface waters, sediments and soils associated with direct sowing soybean cultivation in north pampasic region of Argentina. *Environ Pollut* 156(1):61-6.
- Pieniążek D, Bukowska B, Duda W. 2004. Comparison of the effect of Roundup Ultra 360 SL pesticide and its active compound glyphosate on human erythrocytes. *Pestic Biochem Physiol* 79:58-63.
- Poletta GL, Larriera A, Kleinsorge E, Mudry MD. 2009. Genotoxicity of the herbicide formulation Roundup® (glyphosate) in broad-snouted caiman (*Caiman latirostris*) evidenced by the Comet assay and the Micronucleus test. *Mutat Res* 672(2):95-102.
- Prasad S, Srivastava S, Singh M, Shukla Y. 2009. Clastogenic effects of glyphosate in bone marrow cells of Swiss albino mice. *J Toxicol* doi: 10.1155/2009/308985.
- Quaghebeur D, De Smet B, De Wulf E, Steurbaut W. 2004. Pesticides in rainwater in Flanders, Belgium: results from the monitoring program 1997-2001. *J Environ Monit* 6:182-90.
- Rank J, Jensen A-G, Skov P, Pedersen LH, Jensen K. 1993. Genotoxicity testing of the herbicide Roundup and its active ingredient glyphosate isopropylamine using the mouse bone marrow micronucleus test, Salmonella mutagenicity test, and Allium anaphase-telophase test. *Mutat Res* 300(1):29-36.
- Relyea RA. 2005. The impact of insecticides and herbicides on the biodiversity and productivity of aquatic communities. *Ecol Appl* 15(2):618-27.
- Robinson C. 2010. Argentina's Roundup Human Tragedy. Institute of Science in Society, UK. <http://www.i-sis.org.uk/argentinassRoundupHumanTragedy.php>.
- Romano RM, Romano MA, Bernardi MM, Furtado PV, Oliveira CA. 2010. Prepubertal exposure to commercial formulation of the herbicide glyphosate alters testosterone levels and testicular morphology. *Arch Toxicol* 84:309-317.
- Savitz DA, Arbuckle T, Kaczor D, Curtis KM. 1997. Male pesticide exposure and pregnancy outcome. *Am J Epidemiol* 146(12):1025-36.
- Saxton MA, Morrow EA, Bourbonniere RA, Wilhelm SW. 2011. Glyphosate influence on phytoplankton community structure in Lake Erie. *J Great Lakes Res* 37(4):683-90.
- Schiffman R. 2012. Dow and Monsanto's plan to increase the toxic pesticides sprayed in America's heartland. The Huffington Post, Feb 17. [http://www.huffingtonpost.com/richard-schiffman/dow-and-monsanto-team-up-\\_b\\_1256725.html](http://www.huffingtonpost.com/richard-schiffman/dow-and-monsanto-team-up-_b_1256725.html).
- Schneider MI, Sanchez N, Pineda S, Chi H, Ronco A. 2009. Impact of glyphosate on the development, fertility and demography of *Chrysoperla externa* (Neuroptera: Chrysopidae): Ecological approach. *Chemosphere* 76(10):1451-5.
- Sirinathsinghi E. 2012. Pesticide Illnesses and GM Soybeans - Ban on Aerial Spraying Demanded in Argentina. ISIS Report 18 Jan. [http://www.i-sis.org.uk/Pesticide\\_illnesses\\_and\\_GM\\_soya.php](http://www.i-sis.org.uk/Pesticide_illnesses_and_GM_soya.php).
- Siviková K, Dianovský J. 2006. Cytogenetic effect of technical glyphosate on cultivated bovine peripheral lymphocytes. *Int J Hyg Environ Health* 209(1):15-20.
- Slack RJ, Gronow JR, Voulvoulis N. 2005. Household hazardous waste in municipal landfills; contaminants in leachate. *Sci Total Environ* 337:119-137.
- Slager RE, Poole JA, LeVan TD, Sandler DP, Alavanja MC, Hoppin JA. 2009. Rhinitis associated with pesticide exposure among commercial pesticide applicators in the Agricultural Health Study. *Occup Environ Med* 66(11):718-24.
- Stewart M, Aherns M, Olsen G. 2009. *Field Analysis of Chemicals of Emerging Environmental Concern in Auckland's Aquatic Sediments*. ARC Technical Report 2009/021. Prepared by National Institute of Water and Atmosphere for Auckland Regional Council, Auckland.
- Székács A, Davas B. 2011. Forty years with glyphosate. In: Hasaneen MNAE-G (Ed). *Herbicides - Properties, Synthesis and Control of Weeds*. InTech, Rijeka, Croatia.
- Tenuta M, Beauchamp EG. 1996. Denitrification following herbicide application to a grass sward. *Can J Soil Sci* 76:15-22.
- Trigona M. 2009. Study released in Argentina puts glyphosate under fire. Americas Program Report, July 13. Centre for International Policy, Washington D.C. <http://americas.irc-online.org/am/6254>.
- US EPA. 1992. Pesticides in groundwater database. A compilation of monitoring studies: 1971-1991, National Summary. Office of Prevention, Pesticides and Toxic Substances, United States Environmental Protection Agency, Washington, D.C.
- US EPA. 1993. EPA Reregistration Eligibility Decision Document, Glyphosate. EPA 738-R-93-014. Office of Prevention, Pesticides and Toxic Substances, United States Environmental Protection Agency, Washington, D.C.
- Valente M. 2009. Health Argentina: Scientists reveal effects of glyphosate. 15 Apr, Inter Press Service, Buenos Aires. <http://ipsnews.org>.
- Vera MS, Lagomarsino L, Sylvester M, Pérez GL, Rodríguez P, Mugni H, Sinistro R, Ferraro M, Bonetto C, Zagarese H, Pizarro H. 2009. New evidence of Roundup (glyphosate formulation) impact on periphyton community and the water quality of freshwater ecosystems. *Ecotoxicology* 19(4):710-21.
- Vigfusson NV, Vyse ER. 1980. The effect of the pesticides Dexon, Captan, and Roundup, on sister-chromatid exchanges in human lymphocytes in vitro. *Mut Res* 79:53-7.
- Walsh LP, McCormick C, Martin C, Stocco DM. 2000. Roundup inhibits steroidogenesis by disrupting steroidogenic acute regulatory (StAR) protein expression. *Environ Health Perspect* 108:769-76.
- Wang Y-S, Jaw C-G, Chen Y-L. 1994. Accumulation of 2,4-D and glyphosate in fish and water hyacinth. *Water Air Soil Pollut* 74(3-4):397-403.
- Watts MA. 1994. *The Poisoning of New Zealand*. AIT Press, Auckland.
- WeedScience.com. 2012. International Survey of Herbicide Resistant Plants. <http://www.weedscience.org/In.asp>. Accessed 12 June, 2012.
- Williamson S. 2004. Glyphosate reaps social discontent in South America. *Pestic News* 65:9-10.
- Zablutowicz RM, Reddy KN. 2007. Nitrogenase activity, nitrogen content, and yield responses to glyphosate in glyphosate-resistant soybean. *Crop Prot* 26:370-6.
- Zobiolo LHS, Oliveira RS, Visentainer JV, Kremer RJ, Bellaloui N, Yamada T. 2010. Glyphosate affects seed composition in glyphosate-resistant soybean. *J Agric Food Chem* 58(7):4517-22.